ESPAD

United States ≥ 20 % 15-19 % 10-14 % 5-9 % Non-participating country or data not available Cross-hatching indicates limited comparability or limited geographical coverage.

Daily cigarette use: prevalence in the last 30 days (percentage)



Frequency of alcohol intake in the last 30 days (mean number of occasions among users) United States ≥ 7.0 occasions 6.0-6.9 occasions 5.0-5.9 occasions 4.0-4.9 occasions ≤ 3.9 occasions Non-participating country or data not available Cross-hatching indicates limited geographical coverage.




Average alcohol intake on the last drinking day in centilitres of ethanol among users United States ≥ 6.0 cl 5.0-5.9 cl 4.0-4.9 cl 3.0-3.9 cl ≤ 2.9 cl Non-participating country or data not available Cross-hatching indicates limited comparability or limited geographical



coverage.

Figure 5a. Prevalence of five or more drinks at least once in the last 30 days; one drink contains approximately 2 centilitres of ethanol (percentage)





United States ≥ 14 % 11-14 % 7-10 % 3-6 % ≤ 2 % Non-participating country or data not available Cross-hatching indicates limited comparability or limited geographical coverage.


Prevalence of cannabis use in the last 30 days (percentage) Figure 6a.



Frequency of cannabis use in the last 12 months (mean number of occasions among users) United States ≥ 11.0 occasions 10.0-10.9 occasions 8.0-9.9 occasions 6.0-7.9 occasions ≤ 4.9 occasions Non-participating country or data not available Cross-hatching indicates limited comparability or limited geographical



coverage.



Prevalence of inhalant use in the last 30 days (percentage) Figure 8a.



Figure 9a. Prevalence of new psychoactive substance use in the last 12 months (percentage) United States ≥7% 5-6 % 3-4 % 2 % ≤ 1 % Non-participating country or data not available Cross-hatching indicates limited comparability or limited geographical coverage.



Countries included in the 25-country average (marked in blue) United States

Figure 10.

